

**Section A [Each question carries 1 mark]**

- Charge  $Q$  is kept in a sphere of 5 cm first than it is kept in a cube of side 5 cm. the outgoing flux will be  
 (a) More in case of sphere (c) Same in both case  
 (b) More in case of cube (d) Information Incomplete
- Three capacitors of capacitances  $1\mu\text{f}$ ,  $2\mu\text{F}$  &  $3\mu\text{F}$  are connected in series and a potential difference of 11V is applied across the combination then the potential difference across the plates of  $1\mu\text{f}$  capacitor is  
 (a) 2V (b) 4V (c) 1V (d) 6V
- A wire in the form of a circular loop, of one turn carrying a current, produces magnetic induction  $B$  at the center. If the same wire is looped into a coil of two turns and carries the same current, the new value of magnetic induction at the center is  
 (a)  $B$  (b)  $2B$  (c)  $4B$  (d)  $8B$
- Current sensitivity of a galvanometer can be increased by decreasing:  
 (a) Magnetic field  $B$  (b) number of turns  $N$   
 (c) Torsional constant  $K$  (d) Area  $A$
- The relative permeability of a substance  $X$  is slightly less than unity and that of substance  $Y$  is slightly more than unity, then  
 (a)  $X$  is paramagnetic and  $Y$  is ferromagnetic  
 (c)  $X$  and  $Y$  both are paramagnetic  
 (b)  $X$  is diamagnetic and  $Y$  is ferromagnetic  
 (d)  $X$  is diamagnetic and  $Y$  is paramagnetic
- A wire of magnetic dipole moment  $M$  and  $L$  is bent into shape of a semicircle of radius  $r$ . What will be its new dipole moments?  
 (a)  $M$  (b)  $M/2\pi$  (c)  $M/\pi$  (d)  $2M/\pi$
- A rectangular coil is rotated with a uniform angular velocity about the axis shown in the figure. Initially, the axis of rotation of the coil as well as the magnetic field  $B$  were horizontal. The induced E.M.F. in the coil would be maximum when plane of the coil  
 (a) is horizontal.  
 (b) is at right angle to the magnetic field.  
 (c) makes an angle of  $30^\circ$  with the horizontal.  
 (d) makes an angle of  $45^\circ$  with the direction of magnetic field.
- The magnetic flux through a circuit of resistance  $R$  changes by an amount  $\Delta\phi$  in a time  $\Delta t$ . Total electric charge  $Q$  that passes any point in the circuit during the time  $\Delta t$  is represented by  
 (a)  $Q = \frac{\Delta\phi}{\Delta t}$  (b)  $Q = \frac{\Delta\phi}{R}$  (c)  $Q = R \times \frac{\Delta\phi}{\Delta t}$  (d)  $Q = \frac{1}{R} \times \frac{\Delta\phi}{\Delta t}$
- One requires 11eV of energy to dissociate a carbon monoxide molecule into carbon and oxygen atoms. The minimum frequency of the appropriate electromagnetic radiation to achieve the dissociation lies in  
 (a) visible region (b) infrared region  
 (c) ultraviolet region (d) microwave region
- Which of the following has maximum stopping potential when metal is illuminated by visible light?  
 (a) Blue (b) Yellow (c) Violet (d) Red
- The energy  $E$  of a hydrogen atom with principal quantum no.  $n$  is given by  $E = -13.6/n^2$  eV. The energy ejected when the electron jumps from  $n = 3$  state to  $n = 2$  state of hydrogen is approximately  
 (a) 0.85 eV (b) 1.5 eV (c) 1.9 eV (d) 3.4 eV
- The radius of a nucleus with nucleon number 16 is  $3 \times 10^{-15}$  m. Then, the radius of a nucleus with nucleon number 128 will be :-  
 (a)  $3 \times 10^{-15}$  m (b)  $6 \times 10^{-15}$  m  
 (c)  $9 \times 10^{-15}$  m (d)  $24 \times 10^{-15}$  m

- Assertion (A): The photoelectrons produced by a monochromatic light beam incident on a metal surface, have a spread in their kinetic energies.  
 Reason (R): The work function of the metal varies as a function of depth from the surface.  
 a) Both A and R are true, R is the correct explanation of A.  
 b) Both A and R are true, R is not correct explanation of A.  
 c) A is true but R is false. d) A is false but R is true.

**13. Assertion:-** The electric field at every point is normal to the equipotential surface passing through that point.

**Reason:-** No work is required to move a test charge on an equipotential surface.

- Both Assertion and Reason are correct and the Reason is a correct explanation of the Assertion.
- Both Assertion and Reason are correct but Reason is not a correct explanation of the Assertion.
- The Assertion is correct but Reason is incorrect.
- Both the Assertion and Reason are incorrect.

**14. Assertion :-** When tiny circular obstacle is placed in the path of light from some distance, a bright spot is seen at the centre of the shadow of the obstacle.

**Reason :-** Destructive interference occurs at the centre of the shadow.

- Both Assertion and Reason are correct and the Reason is a correct explanation of the Assertion.
- Both Assertion and Reason are correct but Reason is not a correct explanation of the Assertion.
- The Assertion is correct but Reason is incorrect.
- Both the Assertion and Reason are incorrect.

**15. Assertion :-** Kinetic energy of photo electrons emitted by a photosensitive surface depends upon the intensity of incident photon.

**Reason :-** The ejection of electrons from metallic surface is possible with frequency of incident photon below the threshold frequency.

- Both Assertion and Reason are correct and the Reason is a correct explanation of the Assertion.
- Both Assertion and Reason are correct but Reason is not a correct explanation of the Assertion.
- The Assertion is correct but Reason is incorrect.
- Both the Assertion and Reason are incorrect.


**16. Assertion:-** Silicon is preferred over germanium for making semiconductor devices.

**Reason:-** The energy gap for germanium is more than the energy gap of silicon

- Both Assertion and Reason are correct and the Reason is a correct explanation of the Assertion.
- Both Assertion and Reason are correct but Reason is not a correct explanation of the Assertion.
- The Assertion is correct but Reason is incorrect.
- Both the Assertion and Reason are incorrect.

**Section B [Each question carries 2 marks]**

- Rank the Gaussian surfaces as shown in the figure. In order of increasing electric flux, starting with the most negative



- The refractive index of diamond is much higher than that of glass. How does a diamond cutter make use of this fact?
- Find the radius of curvature of the convex surface of a plano-convex lens, whose focal length is 0.3 m and the refractive index of the material of the lens is 1.5.

(OR)

A telescope consists of two lenses of focal lengths 20 cm and 5 cm. Obtain its magnifying power when the final image is (i) at infinity (ii) at 25 cm from the lenses of eye.

20. If light of wavelength 412.5 nm is incident on each of metals given below, which ones will show photoelectric emission & why

Metal	Work Function (eV)
Na	1.92
K	2.15
Ca	3.20
Mo	4.17

21. Draw the energy band diagram when intrinsic semiconductor (Ge) is doped with impurity atoms of Antimony (Sb). Name the extrinsic semiconductor so obtained and majority charge carriers in it.

(OR)

Draw energy band diagram of p and n type semiconductors.

Also, write two differences between p-type and n-type semiconductors. h to know about the galvanometer before converting it into an ammeter or voltmeter?

**Section C [Each question carries 3 marks ]**


22. A uniformly charged conducting sphere of 2.4 m diameter has a surface charge density of  $80 \mu\text{C}/\text{m}^2$ .

(a) Find the charge on the sphere.

(b) What is the total electric flux leaving the surface of the sphere?

23. Deduce the relationship between current  $I$  flowing through a conductor and drift velocity of the electrons. Following figure shows a plot of current  $I$  flowing through the cross section of a wire versus the time  $T$ .

Use the plot to find the charge flowing in 10 seconds through the wire.




24. (a) What is a moving coil galvanometer?

(b) Give its working principle.


(c) Define the terms: (i) voltage sensitivity and (ii) current sensitivity of a galvanometer.

(OR)

Two parallel straight wires X and Y separated by a distance 5 cm in air carry current of 10 A and 5 A respectively in opposite direction as shown in diagram. Calculate the magnitude and direction of the force on a 20 cm length of the wire Y.



25. A rectangular conductor LMNO is placed in a uniform magnetic field of 0.5 T. The field is directed perpendicular to the plane of the conductor. When the arm MN of length of 20 cm is moved towards left with a velocity of 10 m/s, calculate the emf induced in the arm. Given the resistance of the arm to be 5 ohm (assuming that other arms are of negligible resistance), find the value of the current in the arm.



26. Name the parts of the electromagnetic spectrum which is

(i) suitable for RADAR systems in aircraft navigations.

(ii) used to treat muscular strain.

(iii) used as a diagnostic tool in medicine. Write in brief, how these waves can be produced?

(OR)

(i) Name the EM waves which are used for the treatment of certain forms of cancer. Write their frequency range.

(ii) Thin ozone layer on top of stratosphere is crucial for human survival. Why?

(iii) Why is the amount of the momentum transferred by the EM waves incident on the surface so small?

27. The ground state energy of hydrogen atom is  $-13.6 \text{ eV}$ . If an electron makes a transition from an energy level  $-1.51 \text{ eV}$  to  $-3.4 \text{ eV}$ , calculate the wavelength of the spectral line emitted and name the series of hydrogen spectrum to which it belongs.

28. Calculate the energy released in MeV in the following nuclear reaction:  ${}_{92}\text{U}^{238} \rightarrow {}_{90}\text{Th}^{234} + {}_2\text{He}^4 + \text{Q}$

Mass of  ${}_{92}\text{U}^{238} = 238.05079 \text{ amu}$

Mass of  ${}_{90}\text{Th}^{234} = 234.043630 \text{ amu}$

Mass of  ${}_2\text{He}^4 = 4.002600 \text{ amu}$

$1 \text{ u} = 931 \text{ MeV}$

**Section D [Each question carries 4 marks ]**

29. Case Study: Read the following paragraph & answer the questions.

Two sources of light which continuously emit light waves of same frequency (or wavelength) with a zero or constant phase difference between them, are called coherent sources. Two independent sources of light cannot act as coherent sources, they have to be derived from the same parent source. In Young's double slit experiment, two identical narrow slits  $S_1$  and  $S_2$  are placed symmetrically with respect to narrow slit  $S$  illuminated with monochromatic light. The interference pattern is obtained on an observation screen placed at large distance  $D$  from  $S_1$  &  $S_2$ .

a) Mention any 2 conditions for sustained interference.

b) In the Young's double slit experiment using a monochromatic light of wavelength  $\lambda$ , what is the path difference (in terms of an integer  $n$ ) corresponding to any point having half the peak intensity?

c) Calculate the ratio of the fringe width for bright and dark fringes in YDS experiment.

(OR)

c) In Young's double slit experiment, while using a source of light of wavelength  $4500 \text{ \AA}$ , the fringe width obtained is 0.4 cm. If the distance between the slits and the screen is reduced to half, calculate the new fringe width.

30. Case Study: Read the following paragraph & answer the questions.

A p-n junction is a single crystal of Ge or Si doped in such a manner that one-half portion of it acts as p-type semiconductor and other half functions as n-type semiconductor. As soon as junction is formed, the holes from the p-region diffuse into the n-region and electrons from n-region diffuse into p-region. This results in the development of potential barrier  $V_b$  across the junction which opposes the further diffusion of electrons and holes through the junction. The small region in the vicinity of the junction which is depleted of free charge carriers and has only immobile ions been called the depletion region.

a) Why is germanium preferred over silicon for making semiconductor devices?

b) Which type of biasing results in a very high resistance of a p n junction diode. Draw a diagram showing this bias.

c) How does the width of the depletion region of a pn junction vary, if the reverse bias applied to it decreases.

(OR)

(c) Name the 2 important processes involved in the formation of a p n junction.

